



WHAT TO EXPECT

Your AUCATZYL treatment process

Actor portrayals.

WHAT IS AUCATZYL®

AUCATZYL is a treatment for adults with acute lymphoblastic leukemia. It is used following disease progression while on or after other treatment. AUCATZYL is a medicine made from your own white blood cells, which have been genetically modified to recognize and attack your leukemic cells.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about AUCATZYL?

AUCATZYL may cause side effects that are life-threatening and can lead to death. Call or see your healthcare provider or get emergency help right away if you get any of the following:

- Fever (100.4°F/38°C or higher)
- Severe nausea, vomiting, diarrhea
- Severe headache
- Dizziness or lightheadedness
- Difficulty breathing
- Confusion
- Fast or irregular heartbeat
- Low blood pressure
- Chills/shivering
- Shaking or twitching (tremor)

It is important to tell your healthcare provider that you received AUCATZYL and to show them your AUCATZYL Patient Wallet Card. Your healthcare provider may give you other medicines to treat your side effects.

Before getting AUCATZYL, tell your healthcare provider about all your medical problems, including if you have or have had:

- Neurologic problems (such as seizures, stroke, new or worsening memory loss)
- Lung or breathing problems
- Heart problems
- A recent or active infection
- Past infections which can be reactivated following treatment with AUCATZYL
- Pregnancy, you think you may be pregnant, or plan to become pregnant
- Breastfeeding

Please see additional **Important Safety Information** on the next page and full **Prescribing Information**, including **BOXED WARNING** and Medication Guide.

My procedures and contacts

Partner with your healthcare provider(s) to record the planned dates, times, and locations of the following key procedures.

BONE MARROW ASSESSMENT TO CONFIRM RELAPSE (IF REQUIRED)

Date: _____ Time: _____

Location: _____

HEALTHCARE PROVIDER

Name: _____ Phone number: _____

Email: _____

LEUKAPHERESIS AND IDM SCREENING

Date: _____ Time: _____

Location: _____

APP/NURSE

Name: _____ Phone number: _____

Email: _____

OPTIONAL STABILIZING TREATMENT

Date: _____ Time: _____

Location: _____

APHERESIS CENTER

Name: _____ Phone number: _____

Address: _____

BONE MARROW ASSESSMENT

Date: _____ Time: _____

Location: _____

INFUSION CENTER

Name: _____ Phone number: _____

Address: _____

LYMPHOEDEPLETION

Date: _____ Time: _____

Location: _____

FIRST INFUSION

Date: _____ Time: _____

Location: _____

SECOND INFUSION

Date: _____ Time: _____

Location: _____

AutolusAssist™ is here to help

AutolusAssist can help with your travel, lodging, and meal support, as well as benefits investigations, claims appeals, and financial support. This includes even if you are uninsured or underinsured.



Live support is available each time you call: a Case Manager is ready to answer questions that you may have about insurance, AUCATZYL CAR T-cell treatment, and support resources.

You can call AutolusAssist 24 hours a day, 7 days a week at [\(855\) 288-5227](tel:18552885227).

Visit AutolusAssist.com to learn more and to find resources that can help plan your journey.

Autolus **Assist**

IMPORTANT SAFETY INFORMATION (cont'd)

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, dietary supplements, and herbal supplements.

HOW WILL I RECEIVE AUCATZYL?

- AUCATZYL is made from your own white blood cells, so your blood will be collected by a process called "leukapheresis" (loo-ka-fer-ee-sis), which will concentrate your white blood cells.
- Your blood cells will be sent to a manufacturing center to make your AUCATZYL.
- While waiting for AUCATZYL to be made, you may take other medicines to stabilize your cancer. This is so that your acute lymphoblastic leukemia does not get worse.
- Within 7 days before you start treatment, a sample of your cells will be taken to confirm your disease burden, this will determine which infusion dose you are given first.
- Before you receive AUCATZYL, your healthcare provider will give you chemotherapy to prepare your body.
- When your AUCATZYL is ready, your healthcare provider will give it to you through a catheter (tube) placed into your vein (intravenous infusion).
- After AUCATZYL infusion, you will be monitored daily for at least 7 days after the first infusion.
- You should plan to stay close to a healthcare facility for at least 2 weeks after getting AUCATZYL. Your healthcare provider will help you with any side effects that may occur.
- You may be hospitalized for side effects. Your healthcare provider will determine when you can go home.
- You will receive your second infusion of AUCATZYL 8-12 days after the first infusion.
- Your healthcare provider will need to do blood tests to follow your progress. These blood tests are important. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

WHAT SHOULD I AVOID AFTER RECEIVING AUCATZYL?

- Avoid driving for at least 2 weeks after you get AUCATZYL.
- Do not donate blood, organs, tissues, or cells for transplant.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF AUCATZYL?

- The most common side effects of AUCATZYL include:

<ul style="list-style-type: none"> • Fever (100.4°F/38°C or higher) • Nausea • Diarrhea • Infections • Headache • Muscle or joint pain 	<ul style="list-style-type: none"> • Fatigue or feeling very tired • Low blood pressure (dizziness or lightheadedness, headache, fatigue, short of breath) • Confusion • Difficulty speaking or slurred speech • Low white blood cells (can occur with a fever) • Bleeding
--	--

AUCATZYL may increase your risk of getting cancer, including certain types of blood cancers. Your healthcare provider should monitor you for this.

These are not all the side effects of AUCATZYL. Call your healthcare provider about any side effects that concern you. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full **Prescribing Information**, including **BOXED WARNING** and Medication Guide.

© Autolus, Inc. 2026. AUCATZYL, Autolus, AutolusAssist, and associated logos are trademarks of Autolus Therapeutics. All Rights Reserved. 01/2026-US-AU-0121v2

AUCATZYL treatment plan

BEFORE TREATMENT WITH AUCATZYL



LEUKAPHERESIS AND MANUFACTURING

Your white blood cells will be collected through a process called leukapheresis (sometimes called apheresis). Your blood will also be tested for infectious disease markers (IDMs) to verify that you don't have active viruses.

Q Blood is drawn from a vein in your arm or through a catheter. Your white blood cells are separated and collected, and the remaining blood is returned to your body. The entire process usually takes about 2 to 3 hours. The collected cells are then sent to a facility where they will be modified to make AUCATZYL. It typically takes a few weeks to turn your cells into CAR T cells. In the meantime, you may receive other medicines to treat your cancer.



BONE MARROW ASSESSMENT

Within 7 days before lymphodepleting chemotherapy, a bone marrow sample will be taken.

Q A piece of your bone marrow will be collected through a hollow needle inserted into your hip bone. The results of this test will enable your doctor to tailor the dose of AUCATZYL to your individual disease.



LYMPHODEPLETING CHEMOTHERAPY

Before you receive AUCATZYL, your healthcare provider will give you lymphodepleting chemotherapy for 4 days.

Q Lymphodepletion prepares your body to receive your new CAR T cells by making room for AUCATZYL in your bone marrow. You'll receive 2 drugs during this process: fludarabine and cyclophosphamide.

DURING TREATMENT WITH AUCATZYL



AUCATZYL INFUSION

You'll receive AUCATZYL in 2 separate infusions, about 10 days apart. This may help reduce the possibility of serious side effects, while helping the number of CAR T cells increase after infusion.

Q Your personalized CAR T cells are stored in bags with your name and date of birth. You'll receive your cells through a vein in your arm or a central venous catheter, and each infusion takes about 30 minutes. Avoid driving for at least 2 weeks after you get AUCATZYL.

AUCATZYL DOSING

DOSE 1

3 days after lymphodepleting chemotherapy

Monitoring for side effects begins with Dose 1 and continues for at least 7 days

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, dietary supplements, and herbal supplements.

DOSE 2

DOSE 2

About 10 days after Dose 1




AFTER TREATMENT WITH AUCATZYL



MONITORING

After your first dose, your healthcare team will monitor your recovery daily for at least 7 days.

Q Plan to stay near a healthcare facility for at least 2 weeks after getting AUCATZYL so that your treatment team can continue to monitor your recovery and help you with any side effects. You and your doctor will decide where exactly you will be monitored after treatment. [AutolusAssist](#) can help you with travel, lodging, and meal support during treatment and monitoring, if you qualify for assistance.



MANAGING POSSIBLE SIDE EFFECTS

Common side effects with AUCATZYL include headache, nausea, and tiredness, but more serious side effects are possible. Your healthcare team will watch for and manage any side effects you might experience. Carefully read the Important Safety Information in this brochure so you know the symptoms to watch for after receiving treatment. Make sure you have your treatment team's contact information and know where to report any side effects. Show your wallet card to any healthcare professional you see, so they know you are receiving AUCATZYL.

Learn more at [AUCATZYL.com](#).

B-ALL=B-cell acute lymphoblastic leukemia; CAR=chimeric antigen receptor.
Please see **Important Safety Information** in this brochure and full **Prescribing Information**, including **BOXED WARNING** and Medication Guide.



IMPORTANT SAFETY INFORMATION (cont'd)
Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, dietary supplements, and herbal supplements.

HOW WILL I RECEIVE AUCATZYL?

- AUCATZYL is made from your own white blood cells, so your blood will be collected by a process called "leukapheresis" (loo-ka-huh-re-sis), which will concentrate your white blood cells.
- Your blood cells will be sent to a manufacturing center to make your AUCATZYL.
- While waiting for AUCATZYL to be made, you may take other medicines to stabilize your cancer. This is so that your acute lymphoblastic leukemia does not get worse.
- Within 7 days before you start treatment, a sample of your cells will be taken to confirm your disease burden, this will determine which infusion dose you will receive first.
- Before you receive AUCATZYL, your healthcare provider will give you chemotherapy to prepare your body.
- When your AUCATZYL is ready, your healthcare provider will give it to you through a catheter (tube) placed into your vein (intravenous infusion).
- After AUCATZYL infusion, you will be monitored daily for at least 7 days after the first infusion.
- You should plan to stay close to a healthcare facility for at least 2 weeks after getting AUCATZYL. Your healthcare provider will help you with any side effects that may occur.
- You may be hospitalized for side effects. Your healthcare provider will determine when you can go home.
- You will receive your second infusion of AUCATZYL 8-12 days after the first infusion.
- Your healthcare provider will need to do blood tests to follow your progress. These blood tests are important. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

WHAT SHOULD I AVOID AFTER RECEIVING AUCATZYL?

- Avoid driving for at least 2 weeks after you get AUCATZYL.
- Do not donate blood, organs, tissues, or cells for transplant.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF AUCATZYL?

- The most common side effects of AUCATZYL include:
 - Fever (100.4°F/38°C or higher)
 - Nausea
 - Diarrhea
 - Infections
 - Headache
 - Muscle or joint pain
 - Fatigue (feeling very tired)
 - Low blood pressure (dizziness or lightheadedness)
 - Headache, fatigue, or short of breath
 - Confusion
 - Difficulty speaking or slurred speech
 - Low white blood cells (can occur with a fever)
 - Bleeding

AUCATZYL may increase your risk of getting cancers including certain types of blood cancers. Your healthcare provider should monitor you for this.

These are not all the side effects of AUCATZYL. Tell your healthcare provider about any side effects that concern you. You are encouraged to report any side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full **Prescribing Information**, including **BOXED WARNING** and Medication Guide.

© Autolus, Inc. 2026. AUCATZYL, Autolus, AutolusAssist, and associated logos are trademarks of Autolus Therapeutics. All Rights Reserved. 012026US-AUC-02192

